

University of Dundee

## Rewiring the Immune Response in COVID-19

Chalmers, James D.; Chotirmall, Sanjay H.

*Published in:*  
American Journal of Respiratory and Critical Care Medicine

*DOI:*  
[10.1164/rccm.202007-2934ED](https://doi.org/10.1164/rccm.202007-2934ED)

*Publication date:*  
2020

*Licence:*  
CC BY-NC-ND

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*  
Chalmers, J. D., & Chotirmall, S. H. (2020). Rewiring the Immune Response in COVID-19. *American Journal of Respiratory and Critical Care Medicine*, 202(6), 784-786. <https://doi.org/10.1164/rccm.202007-2934ED>

### General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

paraseptal emphysema. And although neutrophil elastase activity was not measured in this study, it might be speculated that neutrophil elastase might be an important driver of paraseptal emphysema development. A high amount of neutrophil elastase activity has been linked to emphysema in humans and animals (11). Moreover, increased elastase concentrations in sputum are associated with increased risk of exacerbations (12), suggesting that further understanding of the cellular and molecular orchestration leading to emphysema might give insight that ultimately will translate to modifying therapies for many people suffering from COPD.

A question that arises from this work is how paraseptal emphysema develops. The current notion for the pathologic sequence of events from cigarette smoke to centrilobular emphysema development starts with the inflammation of small airway walls occurring in the center of the lobule. This process leads to bronchiolar-alveolar attachments and surrounding alveolar wall destruction. Because paraseptal emphysema occurs adjacent to the pleura and septa and emphysema animal models have marked changes in capillary segments (i.e., a higher number of nonconnecting segments) on the pleural surface (13), disruptions of pulmonary and/or pleural capillaries might also contribute to paraseptal emphysema. It can be speculated that pulmonary perfusion deficiency may lead to misbalanced inflammatory response and tissue damage repair, resulting in paraseptal emphysema.

In conclusion, the authors should be commended for this elegant contribution to pathologic differences in the small airways and cellular composition between paraseptal and centrilobular emphysema. After a long time, this work brings paraseptal emphysema to the center of the stage. ■

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

Alejandro A. Diaz, M.D., M.P.H.  
Harvard Medical School  
Brigham and Women's Hospital  
Boston, Massachusetts

ORCID ID: 0000-0001-5037-2919 (A.A.D.).

## References

1. Diaz AA, Han MK, Come CE, San José Estépar R, Ross JC, Kim V, *et al.* Effect of emphysema on CT scan measures of airway dimensions in smokers. *Chest* 2013;143:687–693.
2. Vasilescu DM, Martinez FJ, Marchetti N, Galbán CJ, Hatt C, Meldrum CA, *et al.* Noninvasive imaging biomarker identifies small airway damage in severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2019;200:575–581.
3. Han MK, Tayob N, Murray S, Woodruff PG, Curtis JL, Kim V, *et al.*; COPDGene and SPIROMICS Investigators. Association between emphysema and chronic obstructive pulmonary disease outcomes in the COPDGene and SPIROMICS cohorts: a *post hoc* analysis of two clinical trials. *Am J Respir Crit Care Med* 2018;198:265–267.
4. Snider GL, Kleinerman J, Thurlbeck WM, Bengali ZK. The definition of emphysema: report of a National Heart, Lung, and Blood Institute, Division of Lung Diseases workshop. *Am Rev Respir Dis* 1985;132:182–185.
5. Heard BE. A pathological study of emphysema of the lungs with chronic bronchitis. *Thorax* 1958;13:136–149.
6. Araki T, Nishino M, Zazueta OE, Gao W, Dupuis J, Okajima Y, *et al.* Paraseptal emphysema: prevalence and distribution on CT and association with interstitial lung abnormalities. *Eur J Radiol* 2015;84:1413–1418.
7. Smith BM, Austin JH, Newell JD, Jr., D'Souza BM, Rozenshtein A, Ahmed F, *et al.* Pulmonary emphysema subtypes on computed tomography: the MESA COPD study. *Am J Med* 2014;127: e7–e23.
8. Castaldi PJ, San José Estépar R, Mendoza CS, Hersh CP, Laird N, Crapo JD, *et al.* Distinct quantitative computed tomography emphysema patterns are associated with physiology and function in smokers. *Am J Respir Crit Care Med* 2013;188:1083–1090.
9. Hobbs BD, Foreman MG, Bowler R, Jacobson F, Make BJ, Castaldi PJ, *et al.*; COPDGene Investigators. Pneumothorax risk factors in smokers with and without chronic obstructive pulmonary disease. *Ann Am Thorac Soc* 2014;11:1387–1394.
10. Tanabe N, Vasilescu DM, Hague CJ, Ikezoe K, Murphy DT, Kirby M, *et al.* Pathological comparisons of paraseptal and centrilobular emphysema in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2020;202:803–811.
11. Ri A, Hagiyaama M, Inoue T, Yoneshige A, Kimura R, Murakami Y, *et al.* Progression of pulmonary emphysema and continued increase in ectodomain shedding of cell adhesion molecule 1 after cessation of cigarette smoke exposure in mice. *Front Cell Dev Biol* 2018;6:52.
12. Shoemark A, Cant E, Carreto L, Smith A, Oriano M, Keir HR, *et al.* A point-of-care neutrophil elastase activity assay identifies bronchiectasis severity, airway infection and risk of exacerbation. *Eur Respir J* 2019;53:1900303.
13. Schraufnagel DE, Schmid A. Capillary structure in elastase-induced emphysema. *Am J Pathol* 1988;130:126–135.

Copyright © 2020 by the American Thoracic Society



## 3 Rewiring the Immune Response in COVID-19

At the time of writing, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic continues globally, with nearly 15 million documented cases and more than 600,000 deaths worldwide (1). Many countries have recently seen falls in the

number of confirmed cases and are beginning to cautiously reopen following lockdown measures, whereas others are experiencing a continued increase, “second waves” of infection, or localized outbreaks following an initial reduction in cases (2, 3).

This pandemic represents the greatest public health, clinical, and scientific challenge of our generation. Containing viral spread has necessitated unprecedented social and economic change as lockdowns only “temporarily” limit morbidity and mortality. Identifying effective therapies and/or a vaccine remain our only long-term solutions. Simply put, research is the only exit strategy (2–5).

©This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).

Originally Published in Press as DOI: 10.1164/rccm.202007-2934ED on August 5, 2020

Great progress has been made in understanding the pathophysiology of coronavirus disease (COVID-19), the disease caused by infection with SARS-CoV-2. Current concepts regard COVID-19 as a biphasic illness: an initial viral phase with suggested mean duration of 7 days followed by a “hyperinflammatory phase” characterized by host-mediated organ damage and what is widely referred to as a “cytokine storm” (6). A subset of patients with COVID-19 display features consistent with hemophagocytic lymphohistiocytosis (HLH), a fulminant hypercytokinemic syndrome associated with multiorgan failure, cytopenias, and abnormal liver function tests (7). There are, however, important differences in the inflammatory profile of HLH and severe COVID-19, suggesting a distinct pathophysiology. The respiratory manifestation of this systemic inflammatory response is acute respiratory distress syndrome (ARDS), and approximately 30% of patients with COVID-19 from the initial series in Wuhan met criteria for ARDS (8). This model of a hyperinflammatory syndrome occurring a week or more after initial infection, leading to ARDS and a need for mechanical ventilation, is further supported by the benefits of dexamethasone in the recently published RECOVERY (Randomized Evaluation of COVID-19 Therapy) trial (9). In this large, open-label, United Kingdom-based trial, dexamethasone significantly reduced COVID-19 mortality, with the largest effect in those requiring mechanical ventilation (9). Most notably, in the supplementary material, patients starting dexamethasone >7 days after symptom onset demonstrated the most significant benefit, whereas those treated in the first 7 days from symptom onset did not, validating the proposed model of a late hyperinflammatory response that can be therapeutically targeted.

In parallel to evidence of excess inflammation, reports of profound viral-induced immunosuppression have emerged. Specifically, deficient production of IFN- $\alpha$  and - $\beta$  in blood is reported in patients with severe disease, whereas a separate study describes deficient IFN- $\gamma$  and TNF- $\alpha$  production (10, 11). Taken together, severe COVID-19 appears to involve a delicate and dynamic balance between excess inflammation and deficient antiinflammatory and antiviral defenses.

The clinical presentation and immunological profile of COVID-19 is therefore unique, and intensive research aimed at characterizing this is ongoing. Most mortality is likely to be driven by host responses rather than the virus, and there is clear evidence of end-organ damage without evidence of viral invasion.

It is a remarkable triumph for clinical and translational research that we know so much about the inflammatory response to SARS-CoV-2 only 6 months after the publication of its gene sequence. Nevertheless, key questions remain about its pathophysiology: which components of the inflammatory response are key “makers” rather than “markers,” and what should be therapeutically targeted (12)? What inflammatory characteristics distinguish between the different stages of disease, and is this useful for risk stratification? Finally, some aspects of the immune response remain less well characterized, and in particular, the contribution of neutrophils to COVID-19 remains unclear (13).

In this issue of the *Journal*, McElvaney and colleagues (pp. 812–821), in a single-center study from Dublin, Ireland, aim to provide fresh insight into the inflammatory response to COVID-19 (14). The study compared 20 patients requiring intensive care admission for severe COVID-19 infection, 20 patients who were hospitalized with stable COVID-19 infection (who did not require

ICU admission), and 15 healthy controls. The study confirmed prior observations with elevated IL-6, IL-1 $\beta$ , soluble TNF receptor-1, and IL-8 in the plasma of severe cases requiring ICU admission, with elevated ratios of IL-6 and IL-1 $\beta$  to the antiinflammatory cytokine IL-10. Metabolic reprogramming of immune cells refers to complex alterations in cell metabolism, typically occurring upon activation by external stimuli such as cytokines, leading to profound change to their function. Neutrophil metabolic reprogramming, or what the authors refer to as “rewiring,” is reported in cystic fibrosis and other conditions in response to inflammatory cytokines and low-level endotoxemia (15). These observations are extended here with increased levels of HIF1- $\alpha$ , PKM2, and cytosolic lactate in neutrophils from patients with COVID-19. Importantly, however, precise mechanisms for these effects remain undefined and are likely to be multifactorial given the combined effects of proinflammatory mediators and hypoxemia in severe COVID-19. Finally, the authors assessed plasma alpha-1 antitrypsin (AAT), which is increased in COVID-19 compared with healthy controls. As IL-6 levels were higher in the more severely ill patients with COVID-19, interestingly, AAT was similarly increased, and the authors demonstrate that positive outcomes were associated with a lower IL-6 relative to AAT. The authors interpret this as meriting a trial of AAT supplementation as AAT modulates inflammatory cytokine production. Nevertheless, because the differences in IL-6:AAT ratios are primarily driven by high IL-6, directly targeting IL-6 represents an equally logical strategy.

This work extends the characterization of COVID-19 by proposing two novel potential targets: neutrophil metabolic reprogramming and a relative AAT deficiency. There are >500 therapeutic trials in COVID-19 worldwide, and as recently reviewed in the *Journal* (12), many duplicate existing trials or have limited mechanistic work underpinning their choice of target or design. Therefore, studies such as that presented here are important to validate potential targets before embarking on therapeutic trials. Demonstrating a mortality benefit with dexamethasone is a significant breakthrough in antiinflammatory therapy for COVID-19; however, the mortality rates of 17.8% (patients not requiring oxygen), 23.3% (patients requiring oxygen) and 29.3% (patients requiring invasive mechanical ventilation) remain high, and novel therapies will be required (9). How should we prioritize? Anti-IL-6 (such as tocilizumab), anti-IL-1 $\beta$  (such as anakinra), anti-TNF (e.g., infliximab), IFN, or exogenous AAT? If the immune system has been rewired, how do we know which wires to cut?

Although many more trials will report in 2020, the efficiency of therapeutic development for COVID-19 will require a means of prioritizing targets or many trials will never complete. In this respect, translational studies such as this, in addition to proof-of-concept phase 2 programs such as ACCORD (Accelerating COVID-19 Research and Development) in the United Kingdom, are the key to rapidly prioritize or discard potential therapies (16). ■

---

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

James D. Chalmers, M.B. Ch.B., Ph.D.  
 Division of Molecular and Clinical Medicine  
 University of Dundee  
 Dundee, United Kingdom

Sanjay H. Chotirmall, M.D., Ph.D.\*  
Lee Kong Chian School of Medicine  
Nanyang Technological University  
Singapore, Singapore

ORCID IDs: 0000-0001-5514-7868 (J.D.C.); 0000-0003-0417-7607 (S.H.C.).

\*S.H.C. is Associate Editor of *AJRCCM*. His participation complies with American Thoracic Society requirements for recusal from review and decisions for authored works.

## References

- European Centre for Disease Prevention and Control. COVID-19 pandemic [accessed 2020 Jul 21]. Available from: <https://www.ecdc.europa.eu/en/covid-19-pandemic>.
- Loneragan M, Chalmers J. Estimates of the ongoing need for social distancing and control measures post-"lockdown" from trajectories of COVID-19 cases and mortality. *Eur Respir J* 2020;56:2001483.
- Niederman MS, Richeldi L, Chotirmall SH, Bai C. Rising to the challenge of COVID-19: advice for pulmonary and critical care and an agenda for research. *Am J Respir Crit Care Med* 2020;201:1019–1022.
- Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, et al.; Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working Group. The effect of control strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study. *Lancet Public Health* 2020;5:e261–e270. [Published erratum appears in *Lancet Public Health* 5:e260.]
- Chotirmall SH, Martinez FJ, Schumacker PT, Cooke CR, Seam N, Brochard L, et al.; The American Thoracic Society Journal Family. Life at the editorial "COVID frontline". *Am J Respir Crit Care Med* 2020;201:1457–1459.
- Wright DJM. Prevention of the cytokine storm in COVID-19. *Lancet Infect Dis* [online ahead of print] 7 May 2020; DOI: 10.1016/S1473-3099(20)30376-5.
- Mehta P, Porter JC, Manson JJ, Isaacs JD, Openshaw PJM, McInnes IB, et al. Therapeutic blockade of granulocyte macrophage colony-stimulating factor in COVID-19-associated hyperinflammation: challenges and opportunities. *Lancet Respir Med* [online ahead of print] 16 Jun 2020; DOI: 10.1016/S2213-2600(20)30267-8.
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; 395:1054–1062. [Published erratum appears in *Lancet* 395:1038.]
- Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, et al.; RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with Covid-19 - preliminary report. *N Engl J Med* [online ahead of print] 17 Jul 2020; DOI: 10.1056/NEJMoa2021436.
- Remy KE, Mazer M, Striker DA, Ellebedy AH, Walton AH, Unsinger J, et al. Severe immunosuppression and not a cytokine storm characterize COVID-19 infections. *JCI Insight* 2020;140329.
- Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Smith N, et al. Impaired type I interferon activity and inflammatory responses in severe COVID-19 patients. *Science* [online ahead of print] 13 Jul 2020; DOI: 10.1126/science.abc6027.
- Mathioudakis AG, Fally M, Hashad R, Knight S, Felton T, Vestbo J. COVID-19 clinical trials: unravelling a methodological Gordian knot. *Am J Respir Crit Care Med* [online ahead of print] 7 Jul 2020; DOI: 10.1164/rccm.202005-1942ED.
- Zuo Y, Yalavarthi S, Shi H, Gockman K, Zuo M, Madison JA, et al. Neutrophil extracellular traps in COVID-19. *JCI Insight* 2020;5:e138999.
- McElvaney OJ, McEvoy NL, McElvaney OF, Carroll TP, Murphy MP, Dunlea DM, et al. Characterization of the inflammatory response to severe COVID-19 illness. *Am J Respir Crit Care Med* 2020;202:812–821.
- McElvaney OJ, Zaslona Z, Becker-Flegler K, Palsson-McDermott EM, Boland F, Gunaratnam C, et al. Specific inhibition of the NLRP3 inflammasome as an antiinflammatory strategy in cystic fibrosis. *Am J Respir Crit Care Med* 2019;200:1381–1391.
- UK Department of Health and Social Care. COVID-19 treatments could be fast-tracked through new national clinical trial initiative. Press release. 2020 Apr 29 [accessed 2020 Jul 20]. Available from: <https://www.gov.uk/government/news/covid-19-treatments-could-be-fast-tracked-through-new-national-clinical-trial-initiative>.

Copyright © 2020 by the American Thoracic Society



## Subclinical Acute Kidney Injury Is Acute Kidney Injury and Should Not Be Ignored

Serum creatinine (SCr) has long been known to be an imperfect biomarker of kidney function (1). Although consensus definitions of acute kidney injury (AKI) using SCr exist, they remain limited (2). More specifically, owing to the nature of renal reserve, it often takes a significant amount of (tubular) injury to translate into a rise in SCr concentrations from baseline (3). Over the last decade, countless investigations have sought to discover and validate biomarkers of AKI that can identify patients who have early kidney injury (4). This concept of elevation in kidney injury biomarkers in the absence of changes in traditional markers (SCr and urine output [UO]) has been dubbed "subclinical AKI" (5). However, given the mounting evidence and

multiple methods of detecting this injury, perhaps it needs a different name because it will not be subclinical for much longer. It is with this backdrop that, in this issue of the *Journal*, Dépret and colleagues (pp. 822–829) investigated plasma proenkephalin A (penKid), a 5-kD stable breakdown product of enkephalins that accumulates in the setting of reduced glomerular filtration (GFR) as a biomarker of subclinical AKI in critically ill patients (6).

However, this new work is not the first to look at subclinical AKI; in 2011, Haase and colleagues published a pooled multicenter analysis of prospective studies focused on NGAL (neutrophil gelatinase-associated lipocalin), an iron-binding protein that is upregulated in the setting of ischemic kidney injury (7). In a compilation of 2,322 critically ill patients, they demonstrated that patients who had elevated NGAL concentrations in the absence of changes in SCr (NGAL[+]/SCr[–]) had similar outcomes compared with those with elevations in SCr in the absence of changes in NGAL (NGAL[–]/SCr[+]). However, both these groups experienced more adverse outcomes (need for renal replacement therapy, longer length of ICU stay, and higher inpatient mortality) compared with those with no elevations in SCr or NGAL.

§This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).

Supported by NIH grant R21DK113420.

Originally Published in Press as DOI: 10.1164/rccm.202006-2239ED on July 8, 2020